

The Republic of the Union of Myanmar
Ministry of Environmental Conservation and Forestry
Forest Department



Study on Sterilization Methods and Shoot Formation of Teak
(*Tectona grandis* L.f.) Tissue Culture



Pyae Pyae Win, Research Assistant-3
Phyu Phyu Hnin, Research Assistant-2
Shwe Lone, Research Assistant-3
Nwe Nwe Win, Research Assistant-3
Ohn Lwin, Professor

December, 2015

**ကျွန်းတစ်သျှူးစတင်မွေးမြူခြင်းအဆင့်တွင် မျိုးပွားတစ်သျှူးများအား ပိုးသန်ခြင်းနည်းစနစ်များကို
လေ့လာခြင်း**

ပြည့်ပြည့်ဝင်း၊ သုတေသနလက်ထောက်-၃
ဖြူဖြူနှင်း၊ သုတေသနလက်ထောက်-၂
ရွှေလုံး၊ သုတေသနလက်ထောက်-၃
နွယ်နွယ်ဝင်း၊ သုတေသနလက်ထောက်-၃
အုန်းလွင်၊ ပါမောက္ခ

စာတမ်းအကျဉ်း

တစ်သျှူးမျိုးပွားခြင်းသည် မိခင်အပင်နှင့် မျိုးရိုးဗီဇတစ်ထပ်တည်း တူညီသည့် အပင်များ ထုတ်လုပ်နိုင်သော၊ စီးပွားဖြစ်ထုတ်လုပ်နိုင်သော နည်းလမ်းတစ်ခုဖြစ်ပါသည်။ သင့်တော်သော ပိုးသတ်ခြင်း နည်းစနစ် မရှိပါက တစ်သျှူးမျိုးပွားရာတွင် မှိုနှင့်ဘက်တီးရီးယား ပိုးမွှားများ ဝင်ရောက်ခြင်းတို့ ဖြစ်စေနိုင်ပါသည်။ ဤစာတမ်းတွင် ပိုးသတ်ခြင်းနည်းစနစ် (၃)မျိုးအား စမ်းသပ် ဆောင်ရွက်ထားပါသည်။ ၎င်းနည်းစနစ်များအနက် စမ်းသပ်မှုနံပါတ်(၃)သည် အကောင်းဆုံး နည်းလမ်းတစ်ခု ဖြစ်ကြောင်းတွေ့ရှိရပါသည်။

Study on Sterilization Methods and Shoot Formation of Teak (*Tectona grandis* L.f.) Tissue Culture

Pyae Pyae Win, Research Assistant-3
Phyu Phyu Hnin, Research Assistant-2
Shwe Lone, Research Assistant-3
Nwe Nwe Win, Research Assistant-3
Ohn Lwin, Professor

Abstract

Tissue culture is an ideal method for production of genetically identical plants and proved to be commercially feasible method. But without proper sterilization method, tissue culture is susceptible to various fungal and bacterial attacks. In this paper, 3 different sterilization methods are trialed. The study suggests sterilization methods: 1) the tips were dipped in Homine , Kasumin for each 5 minutes and HgCl₂ for 3min, and moved to laminar flow for cutting (culture). In the final step, the cut tips are dipped in Clorox and stir the solution containing the tips by hands for about 30 seconds, 2) The tips were washed with tap water for 15 minutes, Homine for 5 minutes, HgCl₂ for 3 minutes and Clorox solution for 2 minutes and washed with DDW, 3) The tips were washed with tap water for 15 minutes , Homine , Kasumin and Ethanol 70% + Tween 20 for 5 mintues. Then they were washed with DDW and were moved to laminar flow for cutting (culture). The cut tips were dipped in HgCl₂ for 1min. In all steps, after treating chemical solutions, the tips were washed with double distilled water (DDW). In trial 1, test tubes subjected to sterilization method S1 gives 7contamination free test tubes, S2 8 tubes and S3 8 tubes, respectively. In trial 2, method S1 gives 6 tubes, S2 7 tubes and S3 9 tubes. In trial 3, method S1 gives 5 tubes, S2 6 tubes and S3 gives 8 tubes. The result shows that sterilization method No.3 is the most appropriate for Teak tissue culture technique. After 30 - 45 days, the aseptic auxiliary shoots were transferred to a fresh medium supplemented with different concentrations of 6-benzyl amino purine (BAP) and Kinetin (KN) treatments.

Contents

| | Page |
|---------------------------|------|
| စာတမ်းအကျဉ်း | i |
| Abstract | ii |
| 1. Introduction | 1 |
| 2. Objectives | 1 |
| 3. Materials and Methods | 1 |
| 4. Results and Discussion | 4 |
| 5. Conclusion | 5 |
| References | |
| Appendix 1 | |
| Appendix 2 | |
| Appendix 3 | |

Study on Sterilization Methods and Shoot Formation of Teak (*Tectona grandis* L.f.) Tissue Culture

1. Introduction

Teak (*Tectona grandis* L.f.) belongs to the family Verbenaceae, which is an economically important, large deciduous, reaching over 30 m in height in favorable conditions. Its uses included flooring and parquet, ship building, furniture-making and in building construction. Teak is grown naturally in India, Myanmar, Thailand and Laos.

Teak is traditionally reproduced through seeds, but in most cases, germination is difficult due to the hard seed coat, low seed quality and late seed production. Also, controlled pollination has turned out to be difficult in teak. Poor germination rate leading to a low production of seedlings further contributes to the paucity of planting material. To enhance teak plantation development, many efforts have been made in the research on improvement of the establishment of mass multiplication.

At present, tissue culture method is proved to be efficient and commercially feasible method for establishment of teak plantations. Tissue culture technique has become an adequate method for producing genetically identical plants on a large scale and in a short time for the plantation establishment. However, there are many varieties of methods in tissue culture and some are found to produce poor shoot proliferation and susceptible to fungal and bacterial attacks. Therefore, it is necessary to experiment which methods are capable of producing maximum number of vigorous plants for a successful plantation.

There are some studies on the micropropagation techniques especially different media for teak tissue culture. In spite of describing the procedure of the sterilization on the explants, there is little or few study on the effect of sterilization methods on the tissue culture propagation of. That is why this research is to show on the effects of different sterilizations on teak tissue culture propagation. This study will intend to support some techniques in the propagation of teak.

In this research paper, we emphasized in finding the most suitable methods of sterilization and culture establishment.

2. Objective

1. To study the effect of different sterilization methods
2. To point out suitable sterilization methods in tissue culture

3. Materials and Methods

3.1 Explant source and sterilization

90 shoot tips and single nodes of young twigs from sexually mature teak plus trees are taken from Forest Research Institute Hedge Garden. Those teak plus trees or clones are taken from Pyi, Paukkaung and Moe Swe and planted in Forest Research Institute Hedge Garden.

The collected sample was washed thrice with tap water to remove the adherent dust particles and other contaminants. As a standard procedure, those shoot tips are sterilized using various methods. In this research paper, we used three different sterilization methods, 10 shoot tips for each method. The experiment was performed three times for each method. The sterilized tips are monitored for infection data and data is recorded every week. (Without hormone media was used as base standard media for all three methods).Fig 1.

3.2. Sterilization Method (1) (S1)

Firstly, the tips are dipped in Homine and stir it using a magnetic stirrer for 5 minutes. Then the tips are washed with DDW (double distilled water). After that, they are dipped again in Kasumin and stir it using a magnetic stirrer for 5 minutes. Then they are washed with DDW. In another step, they are dipped in HgCl₂ and stir it using a magnetic stirrer for 3min and washed again with DDW. After finishing the wash, the tips are moved to laminar flow for cutting (culture). In the final step, the cut tips are dipped in Clorox and stir the solution containing the tips by hands for about 30 seconds. The tips are washed with DDW after wards.Fig 3. Fig 4.

3.3. Sterilization Method (2) (S2)

The tips are washed with tap water for 15 minutes. After that, they are dipped in Homine and stirred using a magnetic stirrer for 5 minutes. Then the tips are washed with double distilled water DDW. In the next step, the tips are sterilized in HgCl₂ solution and stir it using a magnetic stirrer for 3 minutes. Then the tips are washed with DDW. Afterwards, they are dipped in Clorox solution and stirred using a magnetic stirrer for 2 minutes. Then the tips are washed with DDW. Finally, they are sent to laminar flow for cutting (culture) (Fig 3 and Fig 4.).

3.4. Sterilization Method (3) (S3)

In the beginning, the tips are washed with tap water for 15 minutes. Then, the tips are dipped in Homine and stirred the solution using a magnetic stirrer for 5 minutes. Afterwards, the tips are removed from the solution and washed with DDW. For the next step, the tips are dipped in Kasumin and stirred with magnetic stirrer for 5 minutes. Then they are washed with DDW. Next, the tips are dipped in Ethanol 70% + Tween 20 and stirred again using a magnetic stirrer for 5 minutes. After the stirring, the tips are washed with DDW. When above procedures are done, the tips are sent to lamina flow for cutting (culture). There, finally, they are dipped in HgCl₂ and stir it using hand for 1 min. Then wash with double distilled water DDW.Fig 3. Fig 4

3.5. Media and Culture Conditions

The explants were cultured on Murashing and Skoog (MS) medium containing 3% sucrose was used in all the experiments. The pH of the medium was adjusted to 5.8 prior to the addition of 0.6% agar (HiMedia Laboratories Private Limited Mumba, India). Medium was dispensed in 15 ml aliquots into culture tubes (25×150mm) and closed with non-absorbent cotton plugs. The medium was autoclaved at 1.1 kg cm pressure and 121° C temperature for about 15 min (Fig5).

3.6. Culture Establishment

The nodal explants were then singly placed in test tubes (25×150mm) with without hormone media sucrose (3%) and solidified with 0.6% agar was added to the medium to reduce browning. Medium pH was adjusted to 5.8 before autoclaving. After 30 - 45 days, the aseptic auxiliary shoots were transferred to a fresh medium supplemented with different concentrations of 6-benzyl amino purine (BAP) and Kinetin (KN) treatments.



Fig 1

Fig 2

Fig 3

Fig 4



Fig 5

Fig 6

Fig 7

Fig 8



Fig 9

Fig 10

Fig 11

Fig 1. FRI Hedge Garden

Fig 2. Collecting explants from hedge garden

Fig 3. First stage of sterilization; washing with tap water

Fig 4. Sterilization with different sterilization methods using magnetic stirrer

Fig 5. Culture stage in laminar flow

Fig 6. Aseptic shoot tips after cutting

Fig 7. Placing shoot tips in test tubes with media

Fig 8. Placing shoot tips in test tubes with media

Fig 9. Shoot development after 30-35 days

Fig 11. After 45 days, transferred to subculture (multiplication stage)

4. Results and Discussions

4.1 Plant Response on sterilization

Three different sterilization methods (S1, S2, and S3) were performed, using 10 shoots for each method in one time. The shoot tips were put into different test tubes and kept in a separate shelf for each method. The objective of this test is to find the most effective method of sterilization for producing teak shoot tips free from any kind of contamination. The statuses of the sterilized shoots are kept in check and data recorded for once a week for 30 to 45 days. The shoot tips which are found to be attacked by fungi, bacteria or other forms of contamination are recorded as failure.

Analysis of Variance Table for survival rate

| Source | DF | SS | MS | F | P |
|---------------|----|---------|---------|------|---------|
| Rep | 2 | 2.8889 | 1.44444 | | |
| Sterilization | 2 | 8.2222 | 4.11111 | 9.25 | 0.0316 |
| Error | 4 | 1.7778 | 0.44444 | | |
| Total | 8 | 12.8889 | | | |
| Grand Mean | | 7.11111 | | | CV 9.37 |

Table 1. LSD All-Pairwise Comparisons Test of Survival rate for Replication


| Rep | Mean | Homogeneous Groups |
|-----|--------|--------------------|
| 1 | 7.6667 | A |
| 2 | 7.3333 | A |
| 3 | 6.3333 | A |

Alpha 0.05 Standard Error for Comparison 0.5443
 Critical T Value 2.776 Critical Value for Comparison 1.5113
 Error term used: Rep*Sterilization methods, 4 DF There are no significant pairwise differences among the means.

Table 2. LSD All-Pairwise Comparisons Test of Survival rate for Sterilization Methods

| Sterilization Methods | Mean | Homogeneous Groups |
|-----------------------|--------|--------------------|
| S3 | 8.3333 | A |
| S2 | 7.0000 | AB |
| S1 | 6.0000 | B |

Alpha 0.05 Standard Error for Comparison 0.5443
 Critical T Value 2.776 Critical Value for Comparison 1.5113
 Error term used: Rep*Sterilization methods, 4 DF There are 2 groups (A and B) in which the means are not significantly different from one another.



Status of three sterilization methods for three trials

As described above, in trial 1, test tubes subjected to sterilization method S1 gives 7 contamination free test tubes, S2 8 tubes, respectively. In trial 2, method S1 6 tubes, S2 7 tubes and S3 9 tubes. In trial 3, method S1 gives 5 tubes, S2 6 tubes and S3 gives 8 tubes. Therefore, the sterilization method S3 is found to eliminate contamination other than S1 and S2.

5. Conclusion

Sterilization method is the first and foremost step in tissue culture of teak, not to mention it is a vital step. Contamination posts serious obstacles in tissue culture. This research successfully points out an effective sterilization method. No doubt this finding of effective sterilization method is an important finding for further research in tissue culture of teak.

References

D.M.S.H.K.Ranasinghe and G.M. Berlyn · Vegetative Propagation of Teak By Tissue Culture, Department of Forestry and Environmental Science. University of Sri Jayewardenepura School of Forestry and Environmental Studies, Yale University. New Haven, USA , University of Sri Jayewardenepura(1998)

M.L.W.O.M. Karunaratne, S.E. Peries and W.C.P. Egodawatta, Callus Induction and Organogenesis from leaf explants of *Tectona grandis* Scholars Research Library , Annals of Biological Research, 2014,5(4):74-82

Ohn Lwin, Assistant Director , Forest research Institute , December 2009, Comparative Study On Root Development of Teak (*Tectona grandis* L.f.) for Different Propagation Methods (Tissue Culture, Rooted Cutting, Seedling) at Different Stages

R. Srinivasan, G. Ganapathy Selvam, K. Karthikeyan, C. Chandran, S. Kulothungan and C. Govindasamy, In Vitro Propagation of Shoot and Callus Culture of *Tectona grandis* (L.) , are Global Journal of Biotechnology & Biochemistry 7 (1) : 26 -29, 2012

Saw C. Doo, Forest research Institute, February 1983, Develop Teak Tissue Culture Methodology

APPENDIX 1**Analysis of Variance Table for survival rate**

| Source | DF | SS | MS | F | P |
|---------------|-----------|-----------|-----------|----------|----------|
| Rep | 2 | 2.8889 | 1.44444 | | |
| Sterilization | 2 | 8.2222 | 4.11111 | 9.25 | 0.0316 |
| Error | 4 | 1.7778 | 0.44444 | | |
| Total | 8 | 12.8889 | | | |

Grand Mean 7.11111 CV 9.37

Tukey's 1 Degree of Freedom Test for Nonadditivity

| Source | DF | SS | MS | F | P |
|---------------|-----------|-----------|-----------|----------|----------|
| Nonadditivity | 1 | 0.69878 | 0.69878 | 1.94 | 0.2577 |
| Remainder | 3 | 1.07900 | 0.35967 | | |

APPENDIX 2

LSD All-Pairwise Comparisons Test of Survival rate for Replication

| Rep | Mean | Homogeneous Groups |
|------------|-------------|---------------------------|
| 1 | 7.6667 | A |
| 2 | 7.3333 | A |
| 3 | 6.3333 | A |

Alpha 0.05 Standard Error for Comparison 0.5443
 Critical T Value 2.776 Critical Value for Comparison 1.5113
 Error term used: Rep*Sterilization methods, 4 DF There are no significant pairwise differences among the means.

LSD All-Pairwise Comparisons Test of Survival rate for Sterilization Methods

| Sterilization Methods | Mean | Homogeneous Groups |
|------------------------------|-------------|---------------------------|
| S3 | 8.3333 | A |
| S2 | 7.0000 | AB |
| S1 | 6.0000 | B |

Alpha 0.05 Standard Error for Comparison 0.5443
 Critical T Value 2.776 Critical Value for Comparison 1.5113
 Error term used: Rep*Sterilization methods, 4 DF There are 2 groups (A and B) in which the means are not significantly different from one another.

APPENDIX 3**Original Data of three sterilization methods chemical treatments by means of Replications**

| Sterilization Methods | Replications | Survival rate |
|------------------------------|---------------------|----------------------|
| S1 | 1 | 70% |
| | 2 | 60% |
| | 3 | 50% |
| S2 | 1 | 80% |
| | 2 | 70% |
| | 3 | 60% |
| S3 | 1 | 80% |
| | 2 | 90% |
| | 3 | 80% |